

How to Use L^AT_EX 2 _{ε} Class File for MIRU2013

HANAKO GAZO^{1,a)} TARO NINSHIKI^{1,†1,b)} JIRO RIKAI^{2,c)}

1. Introduction

This document describes how to use the class file, `miru2013e.cls`, designed for extended abstract of MIRU2013. The design is based on ASCII Japanese pL^AT_EX 2 _{ε} .

2. Template and How to Typeset a Manuscript

2.1 Language

All manuscripts must be in English or Japanese.

2.2 Paper length

Paper length must be at most 2 pages. Overlength papers will be unaccepted.

2.3 Abstract and keywords

Abstract and keywords must be omitted.

References

- [1] W. Rice, A. C. Wine, and B. D. Grain, diffusion of impurities during epitaxy, Proc. IEEE, vol. 52, no. 3, pp. 284–290, March 1964.
- [2] H. Tong, Nonlinear Time Series: A Dynamical System Approach, J. B. Elsner, ed., Oxford University Press, Oxford, 1990.
- [3] H. K. Hartline, A. B. Smith, and F. Ratliff, Inhibitory interaction in the retina, in Handbook of Sensory Physiology, ed. M. G. F. Fuortes, pp. 381–390, Springer-Verlag, Berlin.
- [4] Y. Yamamoto, S. Machida, and K. Igeta, “Micro-cavity semiconductors with enhanced spontaneous emission,” Proc. 16th European Conf. on Opt. Commun., no. MoF4.6, pp. 3–13, Amsterdam, The Netherlands, Sept. 1990.

¹ Faculty of Engineering, First University, Yamada 1–2–3, Minato-ku, Tokyo, 105–0123 Japan

² R&D Division, Osaka Corporation, Kawada 4–5–6, Suita-shi, 565–0456 Japan

†¹ Presently with Faculty of Information Science, Second University, Sakae 7–8–9, Naka-ku, Nagoya, 460-0008, Japan

a) hanako@gazo.ac.jp

b) taro@ninshiki.co.jp

c) jiro@rikai.co.jp