

How to Use L^AT_EX 2_ε Class File for MIRU2017

Hanako GAZO^{1,a)} Taro NINSHIKI^{2,b)} Jiro RIKAI^{2,c)}

1. Introduction

This is a sample for MIRU2017 papers. The paper is compiled using ASCII Japanese pL^AT_EX 2_ε.

2. How to prepare extended abstract

2.1 Language

Both Japanese and English are acceptable.

2.2 Paper length

Papers are limited to four pages.

2.3 Author name and affiliation

Please list full names, affiliations and email addresses of authors. Do not provide mailing (physical) addresses. The process of selecting oral papers is single blind.

2.4 Abstract and keywords

Abstract and keywords must be omitted.

References

- [1] W. Rice, A. C. Wine, and B. D. Grain, diffusion of impurities during epitaxy, Proc. IEEE, vol. 52, no. 3, pp. 284–290, March 1964.
- [2] H. Tong, Nonlinear Time Series: A Dynamical System Approach, J. B. Elsner, ed., Oxford University Press, Oxford, 1990.
- [3] H. K. Hartline, A. B. Smith, and F. Ratliff, Inhibitory interaction in the retina, in Handbook of Sensory Physiology, ed. M. G. F. Fuortes, pp. 381–390, Springer-Verlag, Berlin.
- [4] Y. Yamamoto, S. Machida, and K. Igeta, “Micro-cavity semiconductors with enhanced spontaneous emission,” Proc. 16th European Conf. on Opt. Commun., no. MoF4.6, pp. 3–13, Amsterdam, The Netherlands, Sept.1990.

¹ First University

² Second University (Presently with Third Corporation)

^{a)} hanako@gazo.ac.jp

^{b)} taro@ninshiki.co.jp

^{c)} jiro@rikai.co.jp